arXiv:1201.2210v2 [math.AP] 17 May 2012

THE SHARP CORNER FORMATION IN 2D EULER DYNAMICS OF PATCHES: INFINITE DOUBLE EXPONENTIAL RATE OF MERGING

SERGEY A. DENISOV

Abstract. For the 2d Euler dynamics of patches, we investigate the convergence to the singular stationary solution in the presence of a regular strain. We prove that the rate of merging can be made double exponential.

1. Introduction and statement of the results

In this paper, we study the 2d Euler dynamics of patches:

$$\dot{\theta} = \nabla\theta \cdot u,\ u = \nabla^{\perp}\Delta^{-1}\theta,\ \theta(z,0) = \theta_0,\ z = (x,y) \in \mathbb{R}^2,\ \nabla^{\perp} = (-\partial_y, \partial_x) \qquad (1)$$

where the θ_0 is the characteristic function of some compact set (a patch). In our case, this compact set will be a centrally symmetric pair of two simply connected domains with rather regular boundaries.

This problem attracted a lot of attention in both physics and mathematical literature in the last several decades and became a classical one. The existence of weak solutions for Euler equation is due to Yudovich [12]; his result, in particular, ensures that the dynamics of patches is well-defined. In [3], Chemin proved that if the boundary of the patch is sufficiently regular then it will retain the same regularity forever; another proof of that fact was given later by Bertozzi and Constantin [1]. We recommend the wonderful books [2, 4] for introduction to the subject and for simplified proofs. For closely related models (e.g., SQG), there were attempts recently to prove that a singularity can occur in finite time (see, e.g., [5] and related [6, 8, 9]).

Although we have global regularity for the 2d Euler dynamics of contours, very little is known about the mechanism leading to "singularity formation". The latter can be described in various ways: growth in time of the curvature, perimeter, etc. In this paper, we focus on one particular geometric characteristic, the distance between two interacting patches, and show that it can decay as double exponential infinitely in time. We also reveal the mechanism that drives this process.

Let $\Omega^- = \{-z, z \in \Omega\}$, i.e., the image of Ω under the central symmetry. The boundary of Ω will be denoted by Γ.

In $\mathbb{R}^2 \sim \mathbb{C}$, we consider the 2d Euler dynamics given by the initial configuration $\Omega(0)\cup\Omega^-(0)$. The areas of patches, the distance between them, the value of vorticity

---

Keywords: Contour dynamics, corner formation, self-similar behavior, double exponential rate of merging

2000 AMS Subject classification: primary 76B47, secondary 76B03.

– all these quantities are of order one as $t = 0$. As time evolves, the Euler evolution deforms $\Omega(0) \cup \Omega^-(0)$ to a new pair $\Omega(t) \cup \Omega^-(t)$ with $|\Omega(t)| = |\Omega(0)|$ because the flow preserves the area and the central symmetry. These new patches will be separated from each other for all times, i.e. $\Omega(t) \cap \Omega^-(t) = \emptyset$, but the question, however, is how small the distance between them can get? The main result of the paper is

Theorem 1.1. *Let positive* δ_1 *be sufficiently small. Then, there is a simply connected domain* $\Omega(0)$ *satisfying* $\mathrm{dist}(\Omega(0), \Omega^-(0)) \sim 1$, *and a time-dependent compactly supported incompressible odd strain*

$$S(z,t) = (P(z,t), Q(z,t))$$

such that

$$\mathrm{dist}(\Omega(t), 0) \sim \mathrm{dist}(\Omega(t), \Omega^-(t)) \lesssim \exp(-e^{\delta_1 t})$$

where $\Omega(t) \cup \Omega^-(t)$ *is the Euler dynamics of* $\Omega(0) \cup \Omega^-(0)$ *in the presence of the strain* $S(z,t)$. *Moreover, for* $S(z,t)$ *we have*

$$\sup_{t \geq 0, z \neq 0} \frac{|S(z,t)|}{|z|} < \infty \tag{2}$$

Remark 1. The corresponding equation for $\theta(z,t) = \chi_{\Omega(t)} + \chi_{\Omega^-(t)}$ is

$$\dot{\theta} = \nabla\theta \cdot (u + S), \ u = \nabla^\perp \Delta^{-1} \theta, \quad \theta(z,0) = \chi_{\Omega(0)} + \chi_{\Omega^-(0)} \tag{3}$$

(compare this to (1)).

Remark 2. As it will be clear from the proof, these contours will touch each other at $t = +\infty$ and the touching point is at the origin. In the local coordinates around the origin the functions parameterizing the contours converge to $\pm|x|$ in a self-similar way which will be described in detail. In the following elementary lemma, we show that under the S–strain alone no point can approach the origin in the rate faster than exponential as long as assumption (2) is made and so it is the nonlinear term $\nabla\theta \cdot u$ in (3) that produces the "double exponentially" fast singularity formation.

Lemma 1.1. *Let* $S(z,t)$ *be an odd vector field that satisfies (2). Consider* $\theta(z,t) = \chi_{\Upsilon(t)}(z)$ *which solves*

$$\dot{\theta} = \nabla\theta \cdot S(z,t), \quad \theta(z,0) = \chi_{\Upsilon(0)}(z) \tag{4}$$

where $\mathrm{dist}(\Upsilon(0), 0) > 0$ *and* $\Upsilon(0)$ *is a compact set. Then,*

$$\mathrm{dist}(\Upsilon(t), 0) \gtrsim e^{-Ct} \tag{5}$$

Proof. This (4) is a transport equation and we have

$$\dot{z} = -S(z,t), \quad z(0) = z_0$$

for the characteristics $z(z_0, t)$. As $S(z,t)$ is odd, $S(0,t) = 0$. Therefore, the origin is a stationary point. The estimate (2) yields

$$|S(z,t)| < C|z|, \quad \forall t \geq 0 \quad \text{and} \quad \forall z \in \mathbb{C}$$

Therefore,

$$|\dot{r}| \leq Cr, \quad r = |z|^2$$

and

$$r(0)e^{-Ct} \leq r(t)$$

This implies (5) as the patch $\Upsilon(t)$ is convected by the flow. □

On the PDE level, we will construct an approximate solution to the 2d Euler dynamics of contours where the error can be interpreted as the strain. It is important to mention here that two centrally symmetric patches can not approach the origin in the rate faster than double exponential even if one places them in the strain S from the theorem 1.1. In other words, the following estimate holds true

$$\mathrm{dist}(\Omega(t),0)\gtrsim \exp(-e^{qt}) \tag{6}$$

with some positive constant q which depends on the initial configuration. This is an immediate corollary of the lemma 8.1 from [2], page 315. The constant δ_1 from the theorem as well as q in (6) can be changed by a simple rescaling (i.e., by multiplying the value of vorticity by a constant) thus the size of δ_1 is small only when compared to the parameters of the problem.

In $\mathbb{R}^2$, in contrast to $\mathbb{T}^2$, the kernel of Δ^{-1} is easier to write and Δ^{-1} can be defined on compactly supported L^1 functions so we will address the problem on the whole plane rather than on $\mathbb{T}^2$. On the 2d torus, similar results hold.

The interaction of two vortices was extensively studied in the physics literature (see, e.g., [10, 11]). For example, the merging mechanism was discussed in [10] where some justifications (both numerical and analytical) were given. In our paper, we provide rigorous analysis of that process and obtain the sharp bounds.

In [7], the authors study an interesting question of the "sharp front" formation. Loosely speaking, the sharp front forms if, for example, two level sets of vorticity, each represented by a smooth time-dependent curve, converge to a fixed smooth arc as $t\to\infty$. Let the "thickness" of the front be denoted by $\delta_{front}(t)$. In [7], the following estimate for 2d Euler dynamics is given (see theorem 3, p. 4312)

$$\delta_{front}(t)>e^{-(At+B)}$$

with constants A and B depend only on the geometry of the front. The scenario considered in our paper is different as the singularity forms at a point.

The idea of the proof comes from the following very natural question. Consider an active-scalar dynamics

$$\dot\theta=\nabla\theta\cdot\nabla^{\perp}(A\theta) \tag{7}$$

where A is the convolution with a kernel $K_A(\xi)$. In some interesting cases, $A=\Delta^{-\alpha}$ with $\alpha>0$ so K_A is positive and smooth near the origin. It also obeys some symmetries inherited from its symbol on the Fourier side, e.g., is radially symmetric. Perhaps, the most interesting cases are $\alpha=1$ (2d Euler), which is treated in this paper, and $\alpha=1/2$ (SQG), which is somewhat less established in the physics literature but still is very interesting mathematically. If one considers the problem (7) with $A=\Delta^{-\alpha}$ on the 2d torus $\mathbb{T}^2=[-\pi,\pi]^2$, then there is a stationary singular weak solution (the author learned about this solution from [9]), a "cross"

$$\theta_s=\chi_E+\chi_{E^-}-\chi_J-\chi_{J^-} \tag{8}$$

where $E=[0,\pi]^2$ and $J=[-\pi,0]\times[0,\pi]$. One can think about two patches touching each other at the origin and each forming the right angle. This picture is also centrally symmetric. Now, the question is: is this configuration stable? In other words, can we perturb these patches a little so that they will converge to the stationary solution at least around the origin? The flow generated by θ_s is hyperbolic and so is unstable. However, a suitably chosen curve placed into the stationary hyperbolic flow will converge to a sharp corner as time evolves. The problem of course is that the actual flow is induced by the patch itself and so it will be changing in time. That suggests that one has to be very careful with the choice of the initial patch to guarantee that this process is self-sustaining. Nevertheless, that seems possible and thus the mechanism of singularity formation through the hyperbolic flow can probably be justified. We do it here by neglecting the smaller order terms. In general, the application of some sort of fixed point argument seems to be needed. Either way, this scenario is a zero probability event (at least the way the proof goes) if the "random" initial condition is chosen. However, if one wants to see merging for a long but fixed time, then this can be achieved for an open set of initial data so from that perspective our construction is realistic.

We will handle the case $\alpha = 1$ only without trying to make the strain smooth. The issue of whether one can choose $S(z,t) \in C^\infty$ or even drop it completely will be addressed in the separate publication and will, perhaps, require the realization of the full fixed-point argument idea explained above. We were able to do that so far only for the model equation where the convolution kernel is a smooth bump.

For $\alpha > 1$, the same argument shows that the merging happens with exponential attraction to the origin. This case is much easier as the convolution kernel is not singular anymore so, for example, one can take the strain exponentially decaying in time.

For $\alpha < 1$, we expect our technique to show that the contours can touch each other in finite time thus proving the blow-up. This, however, will require serious refinement of the method.

There are other stationary singular weak solutions known for 2d Euler dynamics and for other problems in fluid mechanics. It would be interesting to perform analogous stability analysis for each of them with the goal of, e.g., solving the finite time blow up problem.

The structure of the paper is as follows. In the sections 2,3, and 4, we prove some auxiliary results. The section 5 contains the construction of $\Omega(t), \Gamma(t)$ and the proof of the theorem 1.1. In appendix, we find an approximate self-similar solution to the local equation of the curve's evolution.

2. VELOCITY FIELD GENERATED BY THE LIMITING CONFIGURATION

In this section, we consider the velocity field generated by one particular pair of patches which resembles the limiting ($t = \infty$) configuration around the origin of the dynamics described in theorem 1.1. In $\mathbb{R}^2$, take $\theta_s(z) = \chi_E(z) + \chi_{E^-}(z)$, where $E = [0,1]^2$. Recall that in the case of $\mathbb{T}^2$ analogous configuration (8) is a steady state. Let $x, y > 0$, the other cases can be treated using the symmetry of the problem. We have

$$\nabla\Delta^{-1}\theta_s = C\int_{E\cup E^-} \frac{(x-\xi_1, y-\xi_2)}{(x-\xi_1)^2+(y-\xi_2)^2}d\xi_1 d\xi_2$$

Integrating, we have for the first component

$$C\int_0^1 \Big(\log(x^2+(y+\xi_2)^2) - \log(x^2+(y-\xi_2)^2)\Big)d\xi_2 + r_1(x,y)$$

$$= C\int_0^y \log(x^2+\xi^2)d\xi + r_2(x,y)$$

where $r_{1(2)}(x,y)$ are odd and smooth around the origin. Integrating by parts, we get the following expression for the integral

$$y\log(x^2+y^2)-2y+2x\arctan(y/x)$$

By symmetry, for the second component of the gradient we have

$$x\log(x^2+y^2)-2x+2y\arctan(x/y)$$

Thus, the velocity is

$$u_s(z)=\nabla^\perp\Delta^{-1}\theta_s=C(-x,y)\log(x^2+y^2)+r(x,y) \tag{9}$$

around the origin, where $|r(z)|\lesssim|z|$. By scaling the time in Euler dynamics, we can always assume that $C=1/2$ in (9) so we get

Lemma 2.1. *Around the origin, we have*

$$u_s(z)=\nabla^\perp\Lambda_s(z)+r(z) \tag{10}$$

where

$$\Lambda_s(z)=xy\log Q(x,y),\quad Q(x,y)=\begin{cases}|y|, & |y|>|x|\\ |x|, & |x|>|y|\end{cases} \tag{11}$$

and

$$|r(z)|\lesssim|z| \tag{12}$$

Proof. For $x,y>0$, this is a direct corollary of the calculations above. For the other cases, this follows from the symmetry of configuration. $\square$

Remark 1. The function $\nabla^\perp\Lambda_s$ is not continuous on the diagonal. That, however, is not so important as we are trying to control not smoothness of the patches but rather their distance to the origin. This particular choice of Λ_s was made only to simplify the calculations below.

We will focus later on the first term in (10) as the second one has a smaller size and will be absorbed into the strain later (see the formulation of the main theorem). The level sets of Λ_s around the origin are hyperbolas asymptotically. Indeed, take $x=\epsilon\widehat{x}, y=\epsilon\widehat{y}$ and consider

$$\Lambda_s(z)=\epsilon^2\log\epsilon$$

or

$$\widehat{x}\widehat{y}\left(1+\frac{\log Q(\widehat{x},\widehat{y})}{\log\epsilon}\right)=1 \tag{13}$$

For the velocity,

$$\nabla^\perp\Lambda_s(z)=\begin{cases}(-x(\log y+1),y\log y), & y>x\geq 0\\ (-x\log x,y(\log x+1)), & 0\leq y<x\end{cases} \tag{14}$$

For other x and y, we can use the symmetry of Λ_s.

These formulas show that the flow generated by Λ_s is hyperbolic around the origin. Moreover, the attraction and repelling is double exponential. For example, the point $(0,y_0)$ will have a trajectory $(0,y_0^{e^t})$. In the next section, we will study the Cauchy problem associated to this flow.

3. THE CAUCHY PROBLEM FOR THE DYNAMICS GENERATED BY THE LIMITING CONFIGURATION

The construction of the patches and the strain in the main theorem will be based on the calculations done in this section. We consider the following Cauchy problem where t is a parameter:

$$\dot{z}(\tau,t)=\nabla^{\perp}\Lambda_s(z(\tau,t)),\quad \tau\geq 0$$

the formula for the right hand side is given in (14), and the initial position is

$$z(0,t)=(\epsilon(t),e^{-1}),\quad \epsilon(t)=\exp(-e^{\delta t}) \tag{15}$$

For now, δ is some fixed positive number. So, at each time t, a point (let us say it has an **index** t) with initial position $(\epsilon(t),e^{-1})$ starts moving under the flow so that at time $t+\tau$ (i.e., time τ spent since the start of the motion) we will see the arc built by points that are indexed by $t+\tau_1,\tau_1\in[0,\tau]$. Let us call this arc $\Gamma_1(t)$ (rotated by $\pi/4$ in the anticlockwise direction, it will be a part of the $\Gamma(t)=\partial\Omega(t)$ from the main theorem).

Our goal in this section is to study the $\Gamma_1(t)$ around the origin when $t\to+\infty$. This is a straightforward and rather tedious calculation which we have to perform.

For each t, consider the trajectory $z(\tau,t)$ within the fixed neighborhood of the origin: $\{[0,e^{-1}]\times[0,e^{-1}]\}$. This trajectory will be an invariant set for potential Λ_s that corresponds to $\Lambda_s=-e^{-1}\epsilon(t)$. The parameter t, again, can be regarded as the tag of this trajectory. For $t\to\infty$, around zero these trajectories will look more and more like hyperbolas due to (13).

There will be two important events for each trajectory: the first is when it crosses the line $y=x$ at time $\tau=T_1(t)$ and the second is when it crosses the line $x=e^{-1}$ and thus leaves the domain of interest $0<x<e^{-1},0<y<e^{-1}$. We will denote this time by $T_2(t)$. Since the trajectory is an invariant set for Λ_s,

$$z(t,T_1(t))=(\widehat{\epsilon}(t),\widehat{\epsilon}(t))$$

and $\widehat{\epsilon}(t)$ can be found from the equation

$$\widehat{\epsilon}^2(t)\log\widehat{\epsilon}(t)=-e^{-1}\epsilon(t) \tag{16}$$

This gives

$$\widehat{\epsilon}(t)=\sqrt{\frac{2\epsilon(t)}{e\log\epsilon^{-1}(t)}}\left(1+\bar{o}(1)\right),\quad t\to\infty \tag{17}$$

One can write an asymptotical expansion up to any order but we are not going to need it.

Within $y>x>0$, we have $\dot{y}=y\log y$, $y(0,t)=e^{-1}$ and so

$$y(\tau,t)=e^{-e^{\tau}},\quad \tau<T_1(t)$$

Then, the value of $T_1(t)$ can be found from

$$e^{-e^{T_1(t)}}=\widehat{\epsilon}(t)$$

or (due to (16))

$$2e^{T_1}-T_1=1+e^{\delta t}$$

That gives us an asymptotics

$$T_1(t)=\delta t-\log 2+e^{-\delta t}(1-\log 2+\delta t)+O(t^2)e^{-2\delta t} \tag{18}$$

Therefore, at the actual time w,

$$w = t + T_1(t) = t(1+\delta) - \log 2 + e^{-\delta t}(1 - \log 2 + \delta t) + O(t^2)e^{-2\delta t}$$

the curve we study will intersect the line $y = x$ at the point $(\widehat{\epsilon}(t), \widehat{\epsilon}(t))$. Then, assuming that t is a function of w, we have

$$t(w) = \frac{w}{1+\delta} + \frac{\log 2}{1+\delta} - \frac{2^{-\delta_1}(\delta_1 w + 1 - \log 2 + \delta_1 \log 2)}{1+\delta} e^{-\delta_1 w} + O(e^{-1.5\delta_1 w})$$

$$\delta_1 = \delta/(1+\delta) < 1 \tag{19}$$

and

$$f(w) = \widehat{\epsilon}(t(w)) = F(\delta)\exp\left(-e^{\delta_1 w}2^{-1/(\delta+1)} + \frac{\delta_1(\delta_1 - 1)}{2}w\right)(1 + \overline{o}(1)) \tag{20}$$

$$F(\delta) = \exp\left(\frac{\log 2}{2} - \frac{1}{2} + \frac{\delta_1}{2} - \delta_1 \log 2 + \frac{\delta_1^2 \log 2}{2}\right)$$

3.1. **The asymptotical form of the curve within $e^{-1} > y > x > 0$.** We first compute the scaling limit of the curve in the $\left\{|z| < \left(f(w)\right)^{\kappa}\right\}$ neighborhood of zero where κ is any small positive number. Fix the time w and solve the system backward in time using auxiliary functions $\alpha(\tau, w), \beta(\tau, w)$:

$$\dot{\alpha} = \alpha(\log\beta + 1), \quad \dot{\beta} = -\beta\log\beta$$

and $\alpha(0) = f(w), \beta(0) = f(w)$. We get then

$$\beta(\tau, w) = (f(w))^{e^{-\tau}}, \quad \alpha(\tau, t) = e^{\tau}(f(w))^{2-e^{-\tau}} \tag{21}$$

Take T, the moment at which the curve is studied, and let

$$T = w - \tau, \quad \alpha = f(T)\widehat{\alpha}, \quad \beta = f(T)\widehat{\beta}$$

Given $\widehat{\alpha}$ and $\widehat{\beta}$, the parameters w and τ are now the functions of T, $\widehat{\alpha}$, and $\widehat{\beta}$.

We have equations

$$e^{\tau}(f(w))^{2-e^{-\tau}} = f(w-\tau)\widehat{\alpha}, \quad (f(w))^{e^{-\tau}} = f(w-\tau)\widehat{\beta}$$

Denoting $u = \log\widehat{\alpha}, v = \log\widehat{\beta}$, we get

$$\frac{v}{u} = \frac{e^{-\tau} - \dfrac{\log f(w-\tau)}{\log f(w)}}{2 - e^{-\tau} + \dfrac{\tau}{\log f(w)} - \dfrac{\log f(w-\tau)}{\log f(w)}}$$

Now, let $\tau \in [0, C]$ be fixed and take $w \to +\infty$ which is the same then as taking $T \to \infty$. From (20), we immediately get

$$\frac{v}{u} \to \frac{e^{-\tau} - e^{-\delta_1\tau}}{2 - e^{-\tau} - e^{-\delta_1\tau}} = -s_-(\tau, \delta_1), \quad \omega \to 0$$

First, notice that

$$s_-(\tau, \delta_1) > 0$$

and

$$s_-(\tau, \delta_1) \to \omega, \quad \omega = \frac{1-\delta_1}{1+\delta_1} \in (0,1), \quad \tau \to 0$$

Now, we see that, depending on $\tau \geq 0$, we have different zones for x and different asymptotical regimes. For example, if $\tau \to 0$, the limiting curve is

$$\widehat{y} = \widehat{x}^{-\omega} \tag{22}$$

Remark 1. For the distance to the origin we have

$$\min_{\widehat{x}>0}\Big(\widehat{x}^2 + \widehat{x}^{-2\omega}\Big) = \widehat{x}_m^2 + \widehat{x}_m^{-2\omega} < 2$$

where

$$x_m = \omega^{1/(2\omega+2)} < 1$$

Notice now that $\delta_1 \to 0$ as $\delta \to 0$ and

$$\omega \to 1 \quad \text{if} \quad \delta \to 0$$

so in the $\delta \to 0$ limit the limiting shape for $\tau \ll 1$ is hyperbola, which is approximately the invariant set for Λ_s. This makes the perfect sense as the initial point $(e^{-e^{\delta t}}, e^{-1})$ approaches the separatrix OY slower when $\delta \to 0$. Had it been constant in t, the curve would exactly follow the invariant set for Λ_s.

Where precisely on the rescaled space $\widehat{x}, \widehat{y}$ will this uniform convergence take place? We have

$$\alpha(\tau, t) = e^{\tau}\Big(f(T+\tau)\Big)^{2-e^{-\tau}} \sim \Big(f(T)\Big)^{1+\tau_1(T)} \ll f(T)$$

where $\tau_1(T) \to 0$ arbitrarily slowly.

For each $\tau > 0$, there will be its own part of the curve with the corresponding scaling limit given by

$$\widehat{y} = \widehat{x}^{-s_-(\tau,\delta_1)}$$

For this fixed positive τ we can easily identify the domain where this scaling is going to take place. Indeed, at time T, the formula (20) yields

$$f(T) = F(\delta)\exp\left(-e^{\delta_1 T}2^{-1/(\delta+1)} + \frac{\delta_1(\delta_1 - 1)}{2}T\right)(1 + \overline{o}(1)) \tag{23}$$

For the y–coordinate of the point on the curve in the domain represented by τ, (21) gives

$$y \sim \left[\exp\left(-2^{-1/(1+\delta)}e^{\delta_1 T}\right)\right]^{d_1(\tau)}\left[\exp\left(\frac{\delta_1(\delta_1-1)}{2}T\right)\right]^{e^{-\tau}}$$

where $d_1(\tau) = e^{-(1-\delta_1)\tau} < 1$. Thus, we see that $y \sim (f(T))^{d_1(\tau)\pm}$ and for the x–coordinate we have similarly

$$x \sim (f(T))^{l_1(\tau)\pm}, \quad l_1(\tau) = (2 - e^{-\tau})e^{\delta_1\tau} > 1$$

For any fixed $\tau > 0$, we computed the asymptotical shape of the curve. However, we will also need the bounds on the curve for $\tau(T) \to +\infty$. Given arbitrarily small fixed $\mu > 0$, the the part of the curve in the region $y > \Big(f(T)\Big)^{\mu}$ will be contained in the rectangle

$$0 < x < f(T), \quad \Big(f(T)\Big)^{\mu} < y < e^{-1} \tag{24}$$

The first inequality is obvious and can certainly be improved a lot but this is all we will need later on.

3.2. **Asymptotical form of the curve within** $0 < y < x < e^{-1}$**.** This part of the curve is more complicated. To study it, consider equations

$$\dot\alpha(\tau) = -\alpha(\tau)\log\alpha(\tau), \quad \dot\beta(\tau) = \beta(\tau)(\log\alpha(\tau)+1)$$

with initial conditions $\alpha(0) = f(w), \beta(0) = f(w)$. We are interested in the shape of the curve at time $T = w + \tau$.

$$\alpha(\tau) = (f(w))^{e^{-\tau}}, \quad \beta(t) = (f(w))^{2-e^{-\tau}} e^{\tau} \tag{25}$$

We again scale by $f(T)$ as follows

$$\alpha = f(T)\widehat{\alpha}, \quad \beta = f(T)\widehat{\beta}$$

If $u = \log\widehat{\alpha}, v = \log\widehat{\beta}$ then

$$\frac{v}{u} = \frac{\dfrac{\tau}{\log f(w)} + 2 - e^{-\tau} - \dfrac{\log f(w+\tau)}{\log f(w)}}{e^{-\tau} - \dfrac{\log f(w+\tau)}{\log f(w)}}$$

$$\frac{v}{u} \to \frac{2 - e^{-\tau} - e^{\delta_1\tau}}{e^{-\tau} - e^{\delta_1\tau}} = -s_+(\tau,\delta_1) \tag{26}$$

We again have

$$s_+(\tau,\delta_1) \to \omega$$

so the rescaled curve converges to the graph of

$$\widehat{y} = \widehat{x}^{-\omega}, \quad \omega = \frac{1-\delta_1}{1+\delta_1}$$

and this convergence is uniform on $f(T) < x < \Big(f(T)\Big)^{1-\tau_1(T)}$ where $\tau_1(T)$ is positive and tends to zero arbitrary slowly.

For any fixed $\tau \in (0, \tau_{cr})$, we have

$$s_+(\tau,\delta_1) = -\frac{2 - e^{-\tau} - e^{\delta_1\tau}}{e^{-\tau} - e^{\delta_1\tau}} > 0$$

The critical $\tau_{cr}(\delta_1) = \log\xi_{cr}(\delta_1)$ where $\xi_{cr}(\delta_1)$ is the solution of the equation

$$2 - \frac{1}{\xi} - \xi^{\delta_1} = 0, \quad \xi > 1$$

Notice that $\xi_{cr}(\delta_1) \to 1$ as $\delta_1 \to 1$ and $\xi_{cr}(\delta_1) \to +\infty$ as $\delta_1 \to 0$. Consequently, for the critical value

$$\tau_{cr}(\delta_1) \to 0, \quad \delta_1 \to 1$$

and

$$\tau_{cr}(\delta_1) \to +\infty, \quad \delta_1 \to 0$$

As before, for the subcritical value $\tau < \tau_{cr}$, the curve will have a limiting shape given by

$$\widehat{y} = \widehat{x}^{-s_+(\tau,\delta_1)}$$

however $s_+(\tau,\delta_1)$ decays in τ and $s_+(\tau_{cr},\delta_1) = 0$.

The (x,y)–coordinates of part of the curve corresponding to each $\tau < \tau_{cr}$ is easy to find. We again have

$$f(T) = F(\delta)\exp\left(-e^{\delta_1 T}2^{-1/(\delta+1)} + \frac{\delta_1(\delta_1-1)}{2}T\right)(1+\overline{o}(1))$$

and

$$x \sim (f(T))^{d_2(\tau)\pm}, \quad d_2(\tau) = e^{-\tau(1+\delta_1)} < 1$$
$$y \sim (f(T))^{l_2(\tau)\pm}, \quad l_2(\tau) = 2e^{-\delta_1\tau} - e^{-\tau(1+\delta_1)}$$

and $l_2(\tau) > 1$ for $\tau < \tau_{cr}$. In the x–coordinate, the domain corresponding to $\tau < \tau_{cr}$ will be characterized by $f(T) < x < f^{d_2(\tau_{cr}-)}(T)$. Moreover,

$$d_2(\tau_{cr}) \to 0, \quad \delta_1 \to 0$$

What can be said about the curve for the region with $\tau \geq \tau_{cr}$? For the analysis that follows in the next section we will only need very rough bounds.

Starting at time w, the point with coordinates $(f(w), f(w))$ will go along the trajectory which will then cross the line $x = e^{-1}$ in time

$$\tau = \log\log\frac{1}{f(w)} \sim \delta_1 w - \frac{\log 2}{1+\delta} + \bar{o}(1)$$

From (25) we get for $\tau > \tau_{cr}$ and large w

$$\frac{\beta}{\alpha} = e^{\tau}\Bigl(f(w)\Bigr)^{2-2e^{-\tau}} < \Bigl(f(w)\Bigr)^{2-2e^{-\tau_{cr}}} < \Bigl(f(T/(1+\delta_1))\Bigr)^{2-2e^{-\tau_{cr}}}$$

Therefore, part of the curve corresponding to $\tau > \tau_{cr}$ will be contained in the rectangle

$$[\Bigl(f(T)\Bigr)^{d_2(\tau_{cr}-)}, e^{-1}] \times [0, \Bigl(f(T/(1+\delta_1))\Bigr)^{\zeta(\delta_1)}], \quad \zeta(\delta_1) > 0 \qquad (27)$$

3.3. **Self-similar behavior around the origin.** Summarizing the calculations above, we conclude that around the origin the rescaled curve will converge to the graph of the function $\widehat{y} = \widehat{x}^{-\omega}$. How large is the domain of convergence? If the curve is calculated at time T and if $\tau_1(T)$ is arbitrary positive function such that $\tau_1(T) \to 0$ as $T \to \infty$, then this convergence will be valid in the region

$$\Bigl(f(T)\Bigr)^{1+\tau_1(T)} < x < \Bigl(f(T)\Bigr)^{1-\tau_1(T)}$$

Outside this window, we have different scaling limits for different values of the parameter $\tau > 0$.

Remark 2. What is the nature of the scaling law for $\tau \sim 0$ that we have got? Well, the Euler dynamics is defined by the convolution with the kernel

$$K(z) \sim \log|z|$$

and $|\log(s\epsilon)| = |\log\epsilon| + O(1)$ for $s \in [1, M]$ with any fixed M. Thus, the strength of the created hyperbolic flow is more or less the same within any annulus $\epsilon < |z| < M\epsilon$. Outside this annulus, say, for $|z| = \sqrt{\epsilon}$, the size of the kernel is quite different.

Now, assume that we are given the hyperbolic flow in the whole plane defined by the following equations

$$\begin{cases} \dot{x} = y \\ \dot{y} = x \end{cases}$$

Let us find the evolution of the contour $\mathcal{C}(t)$ under this flow which would be self-similar in the sense that $\mathcal{C}(t) = \mathcal{C}(0)e^{-\delta t}$. Assume that $\mathcal{C}(t)$ is given by the graph of the function $y(x,t)$ and then one gets nonhomogeneous Burgers equation for $y(x,t)$

$$y_t(x,t) = -y_x(x,t)y(x,t) + x$$

By our assumptions,

$$y(x,t) = e^{-\delta t}H(e^{\delta t}x)$$

so we have

$$\delta(\xi H'(\xi)-H(\xi))=\xi-H'(\xi)H(\xi) \tag{28}$$

$$H'=\frac{\delta H+\xi}{\delta\xi+H}$$

If $H=\xi z(\xi)$, then

$$|z-1|^A|z+1|^B=C|\xi|^{-1}, A=\frac{1+\delta}{2}, B=\frac{1-\delta}{2}$$

and so

$$|H-\xi|^A|H+\xi|^B=C$$

Due to scaling, we can assume $C=1$. In the coordinates $H-\xi=\beta, H+\xi=\alpha$, we have

$$\beta=\alpha^{-(1-\delta)/(1+\delta)}$$

(compare with (22)).

Is it possible to find the initial data such that the evolution of the curve is self-similar on the larger interval? The answer to this question is yes. For example, in (15), one can take

$$\epsilon(t)=\exp\Bigl(-e^{t^{\tilde\gamma}}\Bigr)$$

with $\tilde\gamma\in(0,1)$. Then, the distance from the curve to the origin will be $\lesssim \exp\Bigl(-e^{t^{\tilde\gamma_1}}\Bigr)$, $\tilde\gamma_1\in(0,1)$ but the self-similar behavior will take place in a wider relative range (but yet not on the whole ball $B_{e^{-1}}(0)$!). This is a general rule: the slower the Cauchy data jumps from one invariant set of Λ_s to another, the more regular the curve is around the origin.

4. Comparison of the velocity fields generated by $\Gamma_1(t)$ and by the limiting configuration.

Let us recall that we denoted by $\Gamma_1(t)$ the arc constructed in the previous section. Let us take $\Gamma_1(t)$ and close it in the smooth and arbitrary way to produce the simply-connected domain $\Omega_1(t)$ (see Figure 1). We will compare now the velocity fields generated by two patches $E\cup E^-$ and $\Omega_1(t)\cup\Omega_1^-(t)$. Recall that

$$\operatorname{dist}\Bigl(\Omega_1(t),0\Bigr)\sim f(t)$$



Figure 1

We will need to use the following lemma in which the picture above will be rotated by $\pi/4$ degrees in the anticlockwise direction. Let us denote by $\Gamma_2(t), \Omega_2(t)$, and E_2 the resulting sets.

Fix t and assume that some arc (let us call it Γ_2 as we will later apply this lemma to the part of $\Gamma_2(t)$) lies above the graph of $|x|$ and below the graph of a certain function $g(x)$ defined on $[-a, a]$ where $a = 1/(e\sqrt{2})$. In this section, we take $\epsilon = f(t)$.

Lemma 4.1. *Rescale $g(x)$ as follows*

$$\widehat{g}(\widehat{x}) = \epsilon^{-1} g(\epsilon \widehat{x}), \quad |\widehat{x}| < a/\epsilon$$

Assume that $\widehat{g}$ satisfies the following properties:

(a) $0 < C_1 < \widehat{g}(\widehat{x}) < C_2$, $\widehat{x} \in [-1, 1]$ *with some absolute constants $C_{1(2)}$.*

(b) *For $|\widehat{x}| > 1$,*

$$|\widehat{x}| \leq \widehat{g} \leq |\widehat{x}| + h(\widehat{x}), \quad h(\widehat{x}) = \overline{o}(|\widehat{x}|), \quad \widehat{x} \to \infty \tag{29}$$

Consider

$$D(z, t) = \nabla^{\perp} \Delta^{-1} (\chi_{\Omega_2(t) \cup \Omega_2^-(t)} - \chi_{E_2 \cup E_2^-})$$

Then, we have the following estimate for $|z| \gtrsim \epsilon$

$$|D(z,t)| \lesssim |z| \left(1 + \int\limits_{1 < |\widehat{x}| < a/\epsilon} \frac{h(\widehat{x}) + h(-\widehat{x})}{|\widehat{x}| \cdot (|\widehat{x} - \widehat{z}_1| + 1)|} d\widehat{x} \right), \quad z = \epsilon \widehat{z}, \, \widehat{z} = (\widehat{z}_1, \widehat{z}_2)$$

Proof. Take $z \in \mathbb{C}^+$. From the central symmetry, we have

$$\nabla^{\perp} \Delta^{-1} (\chi_{\Omega_2 \cup \Omega_2^-} - \chi_{E_2 \cup E_2^-})(0) = 0$$

Now, let us use the following formula

$$\left| \frac{\xi - z}{|\xi - z|^2} - \frac{\xi}{|\xi|^2} \right| = \frac{|z|}{|\xi - z| \cdot |\xi|}$$

to get

$$|D(z,t)| \lesssim |z| \left(1 + \int\limits_{(E_2 \backslash \Omega_2) \cap \{|z| < a/10\}} \frac{d\xi}{|z - \xi| \cdot |\xi|} + \right.$$
$$\left. + \int\limits_{(E_2 \backslash \Omega_2)^- \cap \{|z| < a/10\}} \frac{d\xi}{|z - \xi| \cdot |\xi|} \right)$$

for any $\epsilon \lesssim |z| < a/10$. Scale by ϵ and notice that $|\widehat{z} - \widehat{\xi}| \geq |\widehat{z}_1 - \widehat{\xi}_1|$ so the first integral can be estimated by

$$C \left(1 + \int_{1 < |\widehat{x}| < a/\epsilon} \frac{h(\widehat{x})}{|\widehat{x}| \cdot (|\widehat{x} - \widehat{z}_1| + 1)} d\widehat{x} \right)$$

For the second integral, we have a similar bound with $h(-\widehat{x})$ by symmetry. □

We also immediately get

Corollary 4.1. *If the function h in (29) satisfies*

$$|h(\widehat{x})| \lesssim |\widehat{x}|^{-\gamma}, \quad \gamma > 0 \tag{30}$$

for $|\widehat{x}| < \epsilon^{-\alpha}$, $\alpha \in (0,1)$ and

$$|\ln \epsilon| \max_{\epsilon^{-\alpha} < |\widehat{x}| < a/\epsilon} \left| \frac{h(\widehat{x})}{\widehat{x}} \right| < C \tag{31}$$

then

$$|D(z,t)| \lesssim |z|, \quad |z| \gtrsim \epsilon$$

Proof. From (30)

$$\int_{1<|\widehat{x}|<\epsilon^{-\alpha}} \frac{h(\widehat{x})}{|\widehat{x}| \cdot (|\widehat{x} - \widehat{z}_1| + 1)} d\widehat{x} \lesssim (|\widehat{z}_1| + 1)^{-1}$$

and (31) yields

$$\int_{\epsilon^{-\alpha}<|\widehat{x}|<a/\epsilon} \frac{h(\widehat{x})}{|\widehat{x}| \cdot (|\widehat{x} - \widehat{z}_1| + 1)} d\widehat{x} \lesssim |\ln \epsilon| \max_{\epsilon^{-\alpha}<|\widehat{x}|<a/\epsilon} \left| \frac{h(\widehat{x})}{\widehat{x}} \right| < C$$

□

Now, let us apply this corollary to our situation. For this, we will use the results from the previous section.

Remark 1. Notice that for each δ, the arc $\Gamma_1(t)$ satisfies the conditions of the corollary with some $\alpha(\delta)$. For $|\hat{x}| > \epsilon^{-\alpha}$, we can use (24) and (27) to get

$$\left| \frac{h(\hat{x})}{\hat{x}} \right| < \Big(f(t/(1+\delta_1)) \Big)^{\zeta}, \quad \zeta > 0$$

and

$$\left| \Big(\log f(t) \Big) f(t/(1+\delta_1)) \right| < \Big(f\Big(\frac{t}{1+\delta_1 +} \Big) \Big)^{\zeta} \to 0$$

Thus, we have

$$\sup_{|z| \gtrsim f(t)} \left| \frac{D(z,t)}{z} \right| < C \tag{32}$$

Remark 2. For $|z| \lesssim f(t)$, we can argue as follows. Instead of E above, consider

$$E(t) = E \backslash \{|z| < f(t)\}$$

Then, similarly to (32),

$$\left| \nabla^{\perp} \Delta^{-1} (\chi_{\Omega_2(t) \cup \Omega_2^-(t)} - \chi_{E_2(t) \cup E_2^-(t)}) \right| \lesssim |z|$$

but now this is true for all $z \in B_1(0)$. For $\nabla^{\perp} \Delta^{-1} \chi_{E_2(t)}$, the direct calculation yields

$$\nabla^{\perp} \Delta^{-1} \chi_{E_2(t) \cup E_2^-(t)} = C \nabla^{\perp} \Big(xy \log f(t) \Big) + O(|z|), \quad |z| \lesssim f(t)$$

and so

$$\nabla^{\perp} \Delta^{-1} \chi_{\Omega_2(t) \cup \Omega_2^-(t)} = C \nabla^{\perp} \Big(xy \log f(t) \Big) + O(|z|), \quad |z| \lesssim f(t) \tag{33}$$

where we again can assume that $C = 1$.

5. CONSTRUCTION OF THE VORTEX PATCH DYNAMICS AND PROOF OF THE MAIN THEOREM.

We first construct an incompressible strain Ψ which satisfies the following properties (see Figure 2 for the upper part of the actual picture):

1. Ψ is odd and is compactly supported.

2. Around the points $(4,4)$ and $(-2,2)$ it is the standard hyperbolic time-independent flow (these are the domain D_1 and D_6). At $(-2,2)$, we choose the separatrices to be $y_1=-x$ and $y_2=2+(x+2)$. The flow is attracting along y_2 and is repelling along y_1. To define the curve $\Gamma(t)$ in D_1 we need the following result which will later guarantee necessary initial conditions for the dynamics around $(0,0)$. Let the local coordinates near $(-2,2)$ be denoted by (ξ,η).

Lemma 5.1. *Fix any $\delta>0$ and consider the standard hyperbolic dynamics around the origin*

$$\begin{cases} \dot\xi=\xi, & \xi(0)=\xi_0\\ \dot\eta=-\eta, & \eta(0)=\eta_0\end{cases}$$

Let $G(\xi)=0$ for $\xi\le 0$ and $G(\xi)=\xi^{-1}\exp(-\xi^{-\delta})$ for $\xi>0$. Consider the evolution of the smooth curve $\Gamma(0)=\{(\xi,G(\xi)),|\xi|<1\}$ under this flow. Call it $\Gamma(t)=\{(\xi,G(\xi,t)),|\xi|<1\}$. Then,

$$G(1,t)=e^{-e^{\delta t}} \tag{34}$$

Proof. This is a straightforward calculation. The point (ξ_0,η_0) moves to (ξ_0e^t,η_0e^{-t}) in time t. Thus, the point $(e^{-t},G(e^{-t}))\in\Gamma(0)$ will move to $(1,e^{-e^{\delta t}})$ in time t. $\square$

Remark 1. Notice that the part of $\Gamma(t)$ that belongs to the left half-plane does not change in time. Within the window $|\xi|<1$, the curve $\Gamma(t)$ is always smooth and converges to the coordinate axis.

3. Around the origin (domain D_2), we choose

$$\Psi(z,t)=\nabla^\perp\Lambda_1\left(\frac{y+x}{\sqrt2},\frac{y-x}{\sqrt2},t\right) \tag{35}$$

where

$$\Lambda_1(z,t)=\begin{cases}\Lambda_s(z), & z\notin[-0.01f(t),0.01f(t)]^2\\ xy\log f(t), & z\in[-0.01f(t),0.01f(t)]^2\end{cases}$$

Recall that Λ_s is defined in (11) and $f(t)$ is introduced in (23). We changed the coordinates in (35) as we want to rotate the picture described in section 3 by $\pi/4$ in the positive direction. We also modified the value of the potential in the $0,01f(t)$ neighborhood of the origin to get rid of the artificial singularity generated by the sharp corner in the limiting configuration. Notice now that

$$\Psi(z,t)=\left(\frac{\log f(t)}{2}+O(1)\right)\nabla^\perp(y^2-x^2)$$

in $\{|z|<mf(t)\}$ with any fixed m.

4. Between D_1 and D_2 the potential can be smoothly interpolated.

5. In $D_{3(5)}$, the flow is laminar with direction perpendicular to the black segments and in the north-eastern direction.



Figure 2

6. The potential between zones D_2 and D_3 can be smoothly interpolated, as well as the potential between D_5 and D_6. In the zone D_7, the potential is zero so the curve is frozen. This zone again is interpolated smoothly between D_1 and D_6.

7. In the zone D_4, we construct non-stationary potential in the following way (only in this zone the flow is essentially time-dependent!). We need an argument that allows an interpolation between two laminar flows and guarantees the prescribed evolution of the curve $\Gamma(t)$ in these laminar zones. What we want is to define dynamics in the regions D_3, D_4, D_5 right after the points on the curve leave D_2. We need to define this dynamics in such a way that the motion of $\Gamma(t)$ is localized to these regions and, moreover, that it does not move in D_5. Once again, in D_3 and D_5 we postulate the flow to be laminar and then we want to define it in D_4. We will do that in the local coordinates.

Assume that potential $\Lambda(z) = -y$ in $B = \{z : -1 < x < 0\} \cup \{z : 1 < x < 2\} \sim D_3 \cup D_5$. This potential generates the laminar flow

$$\dot{\theta} = \nabla\theta \cdot \nabla^{\perp}\Lambda$$

where $\nabla^{\perp}\Lambda(z) = (-1, 0)$. We want to define smooth $\Lambda(z,t)$ in $D_4 = \{z : 0 < x < 1\}$ such that the resulting $\Lambda(z,t)$ is smooth globally on $D_3 \cup D_4 \cup D_5$. Moreover, given smooth decaying $\nu(t)$ (e.g., $\nu \in L^1(\mathbb{R}^+)$ is enough for decay condition), we need to define a curve $\Gamma(0) = \{(x, \gamma(x,0))\}$ that evolves under this flow $\Gamma(t) = \{(x, \gamma(x,t))\}$ such that $\gamma(0,t) = \nu(t)$ and $\gamma(1,t) = 0$. This function $\nu(t)$ is determined by $\Gamma(t)$ in the zone D_2 where it approaches the separatrix in the double exponential rate.

To be more precise, ν is proportional to the distance from $\Gamma(t)$ to this separatrix in the area where D_2 and D_3 meet.

We will look for

$$\Lambda(z,t) = -y - g_1(x)g_2(x-t)$$

where $g_{1(2)}$ are smooth. Then, to guarantee the global smoothness, we need $g_1(x) = 0$ around $x = 0$ and $x = 1$. Now, take a point $(0, \nu(T))$ and trace its trajectory for $t > T$. We have

$$x(t,T) = t - T,\ t \in [T, T+1]$$

and

$$y(t,T) = \nu(T) - \int_T^t \Big(g_1'(\tau - T)g_2(\tau - T - \tau) + g_1(\tau - T)g_2'(\tau - T - \tau) \Big) d\tau,\ \ t \in [T, T+1]$$

Since we want $y(T+1, T) = 0$ and g_1 to vanish on the boundary,

$$\nu(T) = g_2'(-T)\int_T^{T+1} g_1(\tau - T)d\tau = g_2'(-T)\int_0^1 g_1(x)dx$$

and this identity should hold for all $T > 0$. Take any g_1 with mean one, this defines g_2 on the negative half-line as long as we set $g_2(-\infty) = 0$. We can continue it now to the whole line in a smooth fashion to have g_2 globally defined. How do we define the initial curve at $t = 0$? We extend smooth $\nu(t)$ to $t \in [-1, 0]$ arbitrarily and apply the procedure explained above to $t \in [-1, \infty)$. The curve that we see at $t = 0$ will be the needed initial value for the dynamics that starts at $t = 0$. It is only left to mention that to localize the picture in the vertical direction we can multiply $\Lambda(z,t)$ be a suitable cut–off in the y direction.

The part of the curve that is in D_5, D_6, D_7, and the north-western part of D_1 is stationary, it does not move at all (this is easy to ensure by making this part of the curve the level set of the stationary potential $\Lambda(z), \Psi = \nabla^{\perp}\Lambda$). For the rest of the curve, it does change in time and the flow is directed along it in the anti-clockwise direction.

Now that the explicit $\Psi(z,t)$ and the curve $\Gamma(t) = \partial\Omega(t)$ evolving under this flow are defined, we are ready to prove theorem 1.1.

Proof. (Theorem 1.1). We have by construction

$$\dot\theta = \nabla\theta \cdot \Psi(z,t)$$

and $\theta(z,t) = \chi_{\Omega(t)}(z) + \chi_{\Omega^-(t)}(z)$. Let us define $S(z,t)$ by

$$S(z,t) = \Psi(z,t) - \nabla^{\perp}\Delta^{-1}\theta$$

Then (32), (33), and (12) give (2) for $S(z,t)$. $\Big($Here we also observed that the contribution from $(\Omega(t) \cup \Omega^-(t)) \backslash B_{0.1a}(0)$ to $\nabla^{\perp}\Delta^{-1}\theta$ is of order $O(|z|)$ around the origin$\Big)$.

Thus, around the origin, the error $S(z,t)$ will in fact be much smaller than $\Psi(z,t)$. It is odd as θ is even and is also divergence free as the difference of two divergence free vector fields. We get

$$\dot\theta = \nabla\theta \cdot \Big(\nabla^{\perp}\Delta^{-1}\theta + S(z,t)\Big) \tag{36}$$

and the theorem is proved as the dynamics of $\Omega(t)$ satisfies

$$\mathrm{dist}(\Omega(t),\Omega^-(t))=2\mathrm{dist}(\Omega(t),0)\sim f(t)$$

For $f(t)$ we have a double exponential decay due to (23).

□

Remark 2. Strictly speaking, the Cauchy data for the evolution in the domain D_2 will not be given by (34) as the flow will distort it when moving between zones D_1 and D_2. However, this leads only to a minor change (a fixed time increment, in fact), so the same argument goes through.

Remark 3. The vector-field Ψ and the strain S we constructed are not even continuous, e.g., Ψ jumps on the lines $y=\pm x$. That, however, is not an issue as the dynamics can still be defined and we are controlling the size of S, not its smoothness. In fact, we could have chosen $\Lambda_s=xy\log(x^2+y^2)$ in (11). That takes care of the regularity issue but the calculations in section 3 would have been much longer.

Remark 4. The parameter δ_1 in the formulation of the theorem 1.1 corresponds to δ_1 in (19) and can be chosen arbitrarily from the interval $(0,1)$. The size of this interval is determined by the parameters of the problem: the value of vorticity, initial size of the patches and their distance to the origin.

6. APPENDIX: APPROXIMATE SELF-SIMILAR SOLUTION TO THE CONTOUR DYNAMICS

In this section, we address the following question: is it possible to construct the approximate solution to the Euler dynamics of patches such that the self-similarity will persist on the larger set? We will do that in a rather artificial way as we already can make the right guess for what the solution should be.

Assume that the boundary of the simply-connected patch is parameterized by $\gamma(s,t)$. Then, the velocity at every point of the contour can be computed by ([5], formula (1))

$$u(\gamma(\xi,t),t)=C\int_0^{2\pi}\log|\gamma(\xi,t)-\gamma(s,t)|\gamma_s'(s,t)ds$$

This is a simple corollary of the Gauss integration formula.

If we have a centrally symmetric pair of vortices interacting with each other and, like before, the part of $\Gamma(t)$ close to the origin can be parameterized by the function $y(x,t)$, then the equation for evolution reads

$$\dot y(x,t)=C\int_{-0.5}^{0.5}(y'(x,t)-y'(\xi,t))\log\left(\frac{(x-\xi)^2+(y(x,t)-y(\xi,t))^2}{(x+\xi)^2+(y(x,t)+y(\xi,t))^2}\right)d\xi+r(x,y,t) \tag{37}$$

where $r(x,y,t)$ is a contribution from those parts of $\Gamma(t)$ and $\Gamma^-(t)$ that are away from the origin. This $r(x,y,t)$ is therefore smooth and $r(0,0,t)=0$ by symmetry. Let us drop $r(x,y,t)$ and try to find an approximate self-similar solution? In other words, we want $y(x,t)=\epsilon(t)\phi(x/\epsilon(t))$ to satisfy (37) up to some smaller order correction. Substitution into (37) gives

$$\frac{\dot\epsilon}{\epsilon}\left(\phi(\widehat{x})-\phi'(\widehat{x})\widehat{x}\right)= \tag{38}$$

$$C\int\limits_{-0.5\epsilon^{-1}}^{0.5\epsilon^{-1}}(\phi'(\widehat{x})-\phi'(\widehat{\xi}))\log\left(\frac{(\widehat{x}-\widehat{\xi})^2+(\phi(\widehat{x})-\phi(\widehat{\xi}))^2}{(\widehat{x}+\widehat{\xi})^2+(\phi(\widehat{x})+\phi(\widehat{\xi}))^2}\right)d\widehat{\xi}+E(\widehat{x},t)$$

where E is an error we will control later on. Let us rewrite the integral as follows (up to a constant multiple)

$$\int_{-0.5\epsilon^{-1}}^{0.5\epsilon^{-1}}(\phi'(\widehat{x})-\phi'(\widehat{\xi}))(\widehat{x}\widehat{\xi}+\phi(\widehat{x})\phi(\widehat{\xi}))K(\widehat{x},\widehat{\xi})d\widehat{\xi}$$

with

$$K(\widehat{x},\widehat{\xi})=\frac{1}{b-a}\int_a^b\frac{d\eta}{\eta},$$
$$a=(\widehat{x}-\widehat{\xi})^2+(\phi(\widehat{x})-\phi(\widehat{\xi}))^2,\quad b=(\widehat{x}+\widehat{\xi})^2+(\phi(\widehat{x})+\phi(\widehat{\xi}))^2$$

and so we get

$$k_1(\widehat{x},t)\phi'(\widehat{x})\widehat{x}+k_2(\widehat{x},t)\phi'(\widehat{x})\phi(\widehat{x})+k_3(\widehat{x},t)\widehat{x}+k_4(\widehat{x},t)\phi(\widehat{x})\tag{39}$$

where the coefficients k_j are defined correspondingly. Assume now that ϕ satisfies the following assumptions:

(a) $\phi(\widehat{x})$ is smooth
(b) $0<C_1<\phi(\widehat{x})<C_2$ for $\widehat{x}\in[-1,1]$
(c) $\phi(\widehat{x})=|\widehat{x}|+\rho(\widehat{x})$ where $|\widehat{x}|>1$ with

$$|\rho(\widehat{x})|<|\widehat{x}|^{-\gamma},\quad|\rho'(\widehat{x})|<|\widehat{x}|^{-\gamma},\quad\gamma>0$$

Let us estimate the coefficients k_j now. We will handle k_3, the analysis for k_2 is similar.

$$k_3(\widehat{x})=-\frac{1}{4}\int_{-0.5\epsilon^{-1}}^{0.5\epsilon^{-1}}\frac{\xi\phi'(\xi)}{\xi\widehat{x}+\phi(\xi)\phi(\widehat{x})}\left(\int_a^b\frac{d\eta}{\eta}\right)d\xi\tag{40}$$

Consider $\widehat{x}\in[1,0.5\epsilon^{-1}]$, the other values can be treated similarly. For the integral over the positive ξ we have (after the change of variables $\xi=\widehat{x}\xi_1$, recall that $x=\epsilon\widehat{x}$)

$$I_1=-\int\limits_0^{0.5x^{-1}}\frac{\xi_1(1+\rho'(\xi_1\widehat{x}))}{\xi_1+(1+\rho(\widehat{x})/\widehat{x})(\xi_1+\rho(\widehat{x}\xi_1)/\widehat{x})}\cdot A\,d\xi_1$$

$$A=\frac{1}{4}\int_{(\xi_1-1)^2+(1-\xi_1+\rho(\widehat{x})/\widehat{x}-\rho(\widehat{x}\xi_1)/\widehat{x})^2}^{(1+\xi_1)^2+(1+\xi_1+\rho(\widehat{x})/\widehat{x}+\rho(\widehat{x}\xi_1)/\widehat{x})^2}\frac{d\eta}{\eta}$$

For A, we have a representation

$$A=\frac{1}{\xi_1}+\frac{\rho(\widehat{x})}{2\widehat{x}\xi_1}+O(\xi_1^{-2}),\quad\xi_1>1$$

so

$$I_1=-\frac{1}{2}\int\limits_1^{0.5x^{-1}}\frac{d\xi_1}{\xi_1}\Big(1+O\left(\frac{\rho(\widehat{x}\xi_1)}{\widehat{x}\xi_1}\right)\Big)\Big(1+\rho'(\xi_1\widehat{x})\Big)+\ldots$$
$$=0.5\log x+O(1)$$

For the other integral, changing the sign in integration

$$I_2 = -\int\limits_{0}^{0.5\epsilon^{-1}} \xi(1+\rho'(-\xi))\frac{1}{b-a}\int_a^b \frac{d\eta}{\eta}d\xi$$

Here, we have

$b = (\widehat{x}-\xi)^2 + (\widehat{x}+\xi+\rho(\widehat{x})+\rho(-\xi))^2,\ a = (\widehat{x}+\xi)^2 + (\widehat{x}-\xi+\rho(\widehat{x})-\rho(-\xi))^2$

As both $\widehat{x}$ and ξ are large in the interesting regime, we are in the situation when

$$a, b > (\widehat{x}^2+\xi^2)/2$$

so we can use the mean-value formula

$$\frac{1}{b-a}\int_a^b \frac{d\eta}{\eta} = \frac{1}{b} + \frac{b-a}{2\eta_1^2}, \quad \eta_1 \in (a,b)$$

Substituting, we have two terms: $I_2 = -(T_1+T_2)$.

$$T_1 = \int_0^{0.5x^{-1}} \xi_1(1+\rho'(-\widehat{x}\xi_1))B^{-1}d\xi_1$$

where

$$B = 2(1+\xi_1^2) + 2(1+\xi_1)(\rho(\widehat{x})/\widehat{x} + \rho(-\widehat{x}\xi_1)/\widehat{x}) \\ +(\rho(\widehat{x})/\widehat{x} + \rho(-\widehat{x}\xi_1)/\widehat{x})^2$$

Thus,

$$T_1 = \int_0^{0.5x^{-1}} \frac{\xi_1(1+\rho'(-\widehat{x}\xi_1))}{2(1+\xi_1^2)}d\xi_1 + O(1) = -0.5\log x + O(1)$$

For the other term, we have

$$|T_2| \lesssim \int_0^{0.5\epsilon^{-1}} \xi\frac{\widehat{x}|\rho(-\xi)| + \xi|\rho(\widehat{x})| + |\rho(\widehat{x})\rho(-\xi)|}{(\widehat{x}^2+\xi^2)^2}d\xi$$

$$\lesssim \int_1^{0.5\epsilon^{-1}} \frac{\xi d\xi}{(\widehat{x}^2+\xi^2)^2}\left(\frac{\widehat{x}}{\xi^\gamma} + \frac{\xi}{\widehat{x}^\gamma} + 1\right)d\xi < C$$

Combining all terms, we have

$$k_3 = \log x + O(1), \quad x > \epsilon$$

For $x \sim 0$, we get $I_{1(2)} = 0.5\log\epsilon + O(1)$. These calculations show that

$$k_3 = \left\{\begin{array}{ll} \log|x| + O(1), & |x| > \epsilon \\ \log\epsilon + O(1), & |x| < \epsilon \end{array}\right. \tag{41}$$

Analogous estimates can be obtained for k_2. They yield

$$k_2 = -\left\{\begin{array}{ll} \log|x| + O(1), & |x| > \epsilon \\ \log\epsilon + O(1), & |x| < \epsilon \end{array}\right. \tag{42}$$

The estimates for other terms are

$$|k_{1(4)}| = O(1)$$

Indeed,

$$k_1 = \int_{-0.5\epsilon^{-1}}^{0.5\epsilon^{-1}} \xi K(\widehat{x},\xi)d\xi, \quad k_4 = -\int_{-0.5\epsilon^{-1}}^{0.5\epsilon^{-1}} \phi'(\xi)\phi(\xi)K(\widehat{x},\xi)d\xi$$

and if one does the same analysis as we did for k_3 in (40), we will get the sum of two integrals: one over positive ξ and the other one over negative ξ. Each will have the same large logarithmic leading term but they will come with different signs now and so will cancel each other in the sum leaving us with the uniformly bounded error terms.

Thus (39) can be written as

$$-\log\epsilon(\phi'\phi-\widehat{x})+E$$

where

$$E=(k_2+\log\epsilon)\phi'\phi+(k_3-\log\epsilon)\widehat{x}+k_1\widehat{x}\phi'(\widehat{x})+k_4\phi(\widehat{x})$$

Going back to (38) and choosing C appropriately ($C<0$), one wants to make the following choice for ϵ and ϕ:

$$\epsilon'=\epsilon\delta_1\log\epsilon,\quad x-\phi'\phi=-\delta_1(\phi-x\phi'),\quad \delta_1\in(0,1)$$

Take for $\epsilon(t)$ one particular solution

$$\epsilon(t)=\exp(-e^{\delta_1 t})$$

The equation for ϕ we had before (see (28)) and so we have

$$|\phi-x|^A|\phi+x|^B=1,\quad A=(1+\delta_1)/2,\quad B=(1-\delta_1)/2$$

and its solution ϕ trivially satisfies assumptions (a), (b), (c) mentioned above.

For the original equation (37), the error one gets after substituting $\epsilon\phi(x/\epsilon)$ amounts to ϵE where

$$\epsilon E\lesssim\epsilon\quad\text{if}\quad |x|\lesssim\epsilon$$

and

$$|\epsilon E|\lesssim\epsilon+\epsilon|(\log|x|-\log\epsilon)(\phi'\phi-\widehat{x})|+|x\phi'(\widehat{x})k_1|+|\epsilon\phi(\widehat{x})k_4|\quad\text{if}\quad |x|\gtrsim\epsilon$$

Therefore, for $|x|\gtrsim\epsilon$,

$$|\epsilon E|\lesssim|z|+\epsilon\left|\frac{x}{\epsilon}\right|^{-\gamma(\delta_1)}\log\left|\frac{x}{\epsilon}\right|,\quad \gamma(\delta_1)>0$$

and $z=(x,\epsilon\phi(\widehat{x}))$. Thus, we see that the error is small again so it is possible to find the approximate solution with the self-similar scaling that holds on the ball of size ~ 1. We are not trying to make this picture global and define the incompressible strain on the whole $\mathbb{R}^2$ which corresponds to the error ϵE but we believe it is possible.

7. Acknowledgment

This research was supported by NSF grant DMS-1067413. The hospitality of the Institute for Advanced Study at Princeton, NJ is gratefully acknowledged. The author thanks A. Kiselev, F. Nazarov, and A. Zlatos for the constant interest in this work and A. Mancho for interesting comments on her preprint [9].

References

[1] A. Bertozzi, P. Constantin, Global regularity for vortex patches, Commun. Math. Physics, 152, (1993), 19–28.

[2] A. Bertozzi, A. Majda, Vorticity and incompressible flow. Cambridge Texts in Applied Mathematics, Cambridge University Press, 2002.

[3] J.-Y. Chemin, Persistence of geometric structures in two-dimensional incompressible fluids, Ann. Sci. Ecole Norm. Sup., (4) 26, (1993), no. 4, 517–542.

[4] J.-Y. Chemin, Perfect incompressible fluids. Oxford Lecture Series in Mathematics and its Applications, 14. The Clarendon Press, Oxford University Press, New York, 1998.

[5] D. Cordoba, M. Fontelos, A. Mancho, J. Rodrigo, Evidence of singularities for a family of contour dynamics equations, PNAS, 102, (2005), no. 17, 5949–5952.

[6] D. Cordoba, On the search for singularities in incompressible flows, Appl. Math., 51, (2006), no. 4, 299–320.

[7] D. Cordoba, C. Fefferman, Behavior of several two-dimensional fluid equations in singular scenarios, Proc. Natl. Acad. Sci. USA, 98, (2001), no. 8, 4311–4312.

[8] D. Cordoba, C. Fefferman, Scalars convected by a two-dimensional incompressible flow, Comm. Pure Appl. Math., 55, (2002), no. 2, 255–260.

[9] A.M. Mancho, Numerical studies on the self-similar collapse of the α-patches problem, preprint, arXiv:0902.0706, (2009).

[10] M.V. Melander, N.J. Zabusky, J.C. McWilliams, Symmetric vortex merger in two dimensions: causes and conditions, J. Fluid Mech., 195, (1988), 303–340.

[11] P.G. Saffman, Vortex dynamics. Cambridge Monographs on Mechanics and Applied Mathematics, Cambridge University Press, 1992.

[12] V.I. Yudovich, Non-stationary flow of an incompressible liquid, Zh. Vychils. Mat. Mat. Fiz., 3, (1963), 1032–1066.

E-mail address: `denissov@math.wisc.edu`

University of Wisconsin-Madison, Mathematics Department, 480 Lincoln Dr. Madison, WI 53706-1388, USA